

ALTERNATIVE TO AMIODARONE IN HEMODYNAMICALLY UNSTABLE VT IV/IO 1.0-1.5 mg/kg Repeat IV/IO 0.5-0.75 mg/kg q 5-10 minutes. Max total dose of 3 mg/kg	
PED: same as adult Adult & PEDs: Repeat boluses I by 50% in pts: 70 and older, pulm edema, hypotensive, liver disease	
ALTERNATIVE TO AMIODARONE IN STABLE VT WITH MD APPROVAL IV/IO 0.50-0.75 mg/kg Repeat IV/IO 0.5-0.75 mg/kg q 5-10 minutes. Maximum total dose of 3 mg/kg	
PED: same as adult Adult & PEDs: Repeat boluses I by 50% in pts: >70, pulmonary edema, hypotensive, liver disease	
LIDOCAINE cont	INTRAOSSEOUS INFUSION FOR CONSCIOUS PTS 20-40 mg of 2 % lidocaine (preservative free) into EZ IO port during intraosseous infusion of fluid bolus or flush
	PED: 0.5 mg/kg of 2 % lidocaine (preservative free) into EZ IO port prior to the initial bolus or flush
MAGNESIUM-	SEIZURES/TOXEMIA IV/IO 4.0 gm of 50% solution diluted with 20 ml NS; IM, 4.0 grams of 50% solution
--SULFATE	V-FIB/PULELESS V-TACH (Bypomagnemia or Torsades) IV/IO 1.0-2.0 gm of 50% solution diluted w/ 20ml NS; PED: IV/IO, 25-50 mg/kg of 50% sol. diluted 5-10 ml NS
NARCAN	OVERDOSE/COMA Adult-IV/IO, IM, SO, 0.4 mg with a dose of 1.6 mg 2 min later if the initial dose is unsuccessful. Up to 5.0 mg may be required. Double the dose for ET admin; 2 mg Intranasally W/MAD device
	IV/IO, IM, SO, 0.1 mg/kg up to a max dose of 2.0 mg. Double dose for ET Rx; MAD device: 0.1mg/kg to max of 2.0 mg
NITROGLYCERIN	CHEST PAIN, PULM EDEMA SUSPECTED ANGINA SL 0.4 mg q 5 min up to 3 tablets as long as BP remains over 100 mm/Hg systolic
OXYMETAZOLINE	EPISTAXIS, NASOTRACHEAL INTUBATION 2-3 sprays in each bleeding nostril or nostril selected for nasotracheal intubation
RACEMIC EPI	STRIDOR > 2 YR: 0.5 ml (2.25%) diluted in 3ml NS via nebulizer; < 2 YR: 0.25ml (2.25%) diluted in 3ml NS via nebulizer
SODIUM-	POST-RESUSCITATION HYPOTENSION IV/IO 1.0 mEq/kg Repeat q 10 minutes with IV/IO 0.5 mEq/kg PED: same as adult
--BICARBONATE	CYCLIC ANTI-DEPRESSANT OD IFORS duration > 100 msec, then IV/IO bolus 1.0 mEq/kg Repeat q 10 minutes X 3; PED: same as adult
	IV/IO 1.0 mEq/kg Repeat q 10 minutes with IV/IO 0.5 mEq/kg PED: same as adult
SUCCINYL-	CRUSH INJURY IV/IO 1.0 mEq/kg HYPERKALEMIA: Add NaHCO3 1.0 mEq/kg to 1000 cc NS wide Open PED: same as adult
--CHOLINE	RSI Pretrat Adult & PEDs with Lidocaine IV/IO 1.0 mg/kg up to 100 mg if heard injury or 2 nd dose of succinylcholine needed
	Pretrat Adults with Atropine 1.5 IV/IO mg prior to 2 nd dose of succinylcholine
	Pretrat conscious pt with Versed or if the patient is hypotensive and older than 10 years, administer Etomidate
	SUX: ADULT or PED: IV/IO 1.5 mg/kg; may repeat once; IM, 3 mg/kg to total of 150 mg
VECURONIUM	MAINTAIN PARALYSIS OF INTUBATED PATIENTS IV/IO 0.1 mg/kg. May repeat x 1. PED: same as adult, do not give if < 7 weeks old
	ALTERNATIVE TO SUCCINYLCHOLINE IV/IO 0.1 mg/kg. May repeat x 1. PED: same as adult, do not give if < 7 weeks old
	Note: Reconstitute powder with 10 ml of Normal Saline (1 ml of NS, to each 1 mg of drug)
VENTOLIN	ASTHMA, COPD, & TOXIC SMOKE INHALATION 2.5 mg in 3 ml NS @ 6L. May repeat PEDs: > 2 YR: 2.5 mg in 3 ml NS @ 6L; < 2 YR: 1.25 mg in 1 ml NS @ 6L
	RENAL FAILURE & CRUSH INJURY SYNDROME 5 mg in 6 ml NS @ 6L. May repeat PEDs: All ages, 2.5 mg in 3 ml NS @ 6L.
VERSED	CARDIOVERSION, ADULT/PED SEIZURES* & Adult: IV/IO, IM 0.5-1.0 mg q 3 min up to a max of 5.0 mg.
	SEDTATION FOLLOWING INTUBATION* PED: IV/IO, IM, PR 0.05-0.1 mg/kg q 3 min to a max dose of 2.0 mg
	PREMEDICATION TO INTUBATION & EXT PACING IV/IO, 2.5-5.0 mg PED: IV/IO, IM, PR 0.05-0.1 mg/kg q 3 min to a max dose of 2.0 mg
	EXCITED DELIRIUM* & SEVERE AGITATION* IV/IO, 2.5 mg q 3-5 min up to max of 10 mg PED: IV/IO, IM, PR 0.05-0.1 mg/kg q 3 min to a max dose of 2.0 mg
	ALL ASTERISKED ADULT & ALL PED INDICATIONS NOTE: *May be given via MAD to adults at dose of 5.0 mg. May be given via MAD to PEDs at 0.2 mg/kg to max of 2.0 mg
ZOFAN	NAUSEA & VOMITING IV/IO, IM 4 mg undiluted, give over 30-300 seconds, may repeat x 1 PED: same as adult for children > 40 kg; 0.1 mg/kg for children 40 kg or less
	DEACONESS 473-7196 (DEDICATED EMS GROUND LINE)
	SACRED HEART 458-7100 (ER DEPT) 474-4840 (DEDICATED EMS GROUND LINE)
	HOLY FAMILY 455-3344 (ER DEPT) 482-2460 (ER DEPT) NO DEDICATED EMS GROUND LINE
	VALLEY GENERAL 473-5466 (ER DEPT) 922-9495 (DEDICATED EMS GROUND LINE)
	VETERANS 434-7000 (ER DEPT) 434-7010 (DEDICATED EMS GROUND LINE)
	POISON CENTER 1-800-7450 M-F 8AM-5PM 1-800-222-1222 (NATIONWIDE)
	ELDER SERVICES 458-7450 M-F 8AM-5PM
	CPS 363-3333 M-F 8AM-5PM 1-800-562-5624 (AFTER HOURS-24/7) DOMESTIC VIOLENCE (326-2255) 24/7 HOTLINE
	SPOKANE MENTAL HEALTH 838-4651 (24/7) FOR CHILD & FAMILY SERVICES CALL 458-7458 M-F 8AM-5PM
	FIRST CALL FOR HELP 838-4428 (24/7 REFERRAL SERVICE) TOLL FREE @ 1-877-678-4428